



THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

Uranium Nitrogen Multiple Bonding: Isostructural Anionic, Neutral, and Cationic Uranium Nitride Complexes Featuring a Linear $\text{U}=\text{N}=\text{U}$ Core

Citation for published version:

Fox, AR, Arnold, PL & Cummins, CC 2010, 'Uranium Nitrogen Multiple Bonding: Isostructural Anionic, Neutral, and Cationic Uranium Nitride Complexes Featuring a Linear $\text{U}=\text{N}=\text{U}$ Core', *Journal of the American Chemical Society*, vol. 132, no. 10, pp. 3250-3253. <https://doi.org/10.1021/ja910364u>

Digital Object Identifier (DOI):

[10.1021/ja910364u](https://doi.org/10.1021/ja910364u)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of the American Chemical Society

Publisher Rights Statement:

Copyright © 2010 by the American Chemical Society. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Journal of the American Chemical Society*, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <http://dx.doi.org/10.1021/ja910364u>

Cite as:

Fox, A. R., Arnold, P. L., & Cummins, C. C. (2010). Uranium Nitrogen Multiple Bonding: Isostructural Anionic, Neutral, and Cationic Uranium Nitride Complexes Featuring a Linear U=N=U Core. *Journal of the American Chemical Society*, 132(10), 3250-3253.

Manuscript received: 08/12/2009; Article published: 24/02/2010

Uranium–Nitrogen Multiple Bonding: Isostructural Anionic, Neutral, and Cationic Uranium Nitride Complexes Featuring a Linear U=N=U Core**

Alexander R. Fox,¹ Polly L. Arnold² and Christopher C. Cummins^{1,*}

^[1]Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Room 6-435, Cambridge, Massachusetts 02139, USA.

^[2]EaStCHEM, School of Chemistry, Joseph Black Building, University of Edinburgh, West Mains Road, Edinburgh, EH9 3JJ, UK.


^[*]Corresponding author; e-mail: ccummins@mit.edu

^[**]We thank the National Science Foundation for supporting this research (CHE-0724158). A.R.F. thanks MIT for the Alan Davison Graduate Fellowship.

Supporting information:

Experimental details, including treatment of $[3][B(Ar^F)_4]$ with CO and LiCCPh; crystallographic data and files in CIF format; and computational data relevant to the electronic structure of **3**. This material is available free of charge via the Internet at <http://pubs.acs.org>

Graphical abstract:



Abstract

Reaction of the uranium(III) tris(anilide) complex (THF)U(N[*t*-Bu]Ar)₃ (**1**, THF = tetrahydrofuran; Ar = 3,5-Me₂C₆H₃) with MN₃ (M = Na, [N(*n*-Bu)₄]) results in the formation of the bimetallic diuranium(IV/IV) complexes M[(μ-N)(U(N[*t*-Bu]Ar)₃)₂] (M[**3**]), which feature a single nitride ligand engaged as a linear, symmetric bridge between two uranium centers. The stability of the U=N=U core across multiple charge states is illustrated by stepwise chemical oxidation of Na[**3**] to the diuranium(IV/V) complex (μ-N)(U(N[*t*-Bu]Ar)₃)₂ (**3**) and the diuranium(V/V) complex [(μ-N)(U(N[*t*-Bu]Ar)₃)₂][B(Ar^F)₄] {[**3**][B(Ar^F)₄]; Ar^F = 3,5-(CF₃)₂C₆H₃}. M[**3**], **3**, and [**3**][B(Ar^F)₄] were characterized by NMR spectroscopy, single-crystal X-ray diffraction, and elemental analysis. The cyclic voltammogram of **3** reveals two clean, reversible one-electron electrochemical events at $E_{1/2} = -1.69$ and -0.67 V, assigned to the [**3**][−]/**3** and **3**/[**3**]⁺ redox couples, respectively. The X-ray crystal structures of [N(*n*-Bu)₄][**3**], **3**, and [**3**][B(Ar^F)₄] reveal a linear U=N=U core that contracts by only ~0.03 Å across the [**3**]^{*n*} (*n* = −1, 0, +1) series, an effect that is rationalized as being primarily electrostatic in origin. [**3**][B(Ar^F)₄] reacts with NaCN, eliminating Na[B(Ar^F)₄] and forming the known diuranium(IV/IV) cyanoimide complex (μ-NCN)(U(N[*t*-Bu]Ar)₃)₂, suggesting that the U=N=U core has metallonitrene-like character.

Introduction

Uranium nitrides have been identified as worthy candidates for N-atom transfer reactions and aziridination of small molecules, and their potential use in catalytic cycles and as advanced nuclear fuels has been noted.¹ To date, only a handful of uranium nitride complexes have been reported, most of which are polymetallic nitride clusters derived from dinitrogen or azide activation² and binary and ternary nitrides generated and observed under matrix-isolation conditions.³ Beyond uranium nitrides, assessing the degree of covalency and mixing of 5f and 6d orbitals in actinide–ligand bonding continues to be an active area of research⁴ and has been recently extended to multiply bonded uranium imido complexes shown to accommodate multiple formal oxidation states at uranium.⁵

We recently reported on the reaction of (THF)U(N[*t*-Bu]Ar)₃ (**1**, THF = tetrahydrofuran; Ar = 3,5-Me₂C₆H₃) with [N(*n*-Bu)₄][B(N₃)(C₆F₅)₃], which directly furnishes the monometallic uranium(V) nitridoborate complex [N(*n*-Bu)₄][(C₆F₅)₃B–N≡U(N[*t*-Bu]Ar)₃] ([N(*n*-Bu)₄][**2**]) and provides access to the neutral uranium(VI) nitridoborate derivative (C₆F₅)₃B–N≡U(N[*t*-Bu]Ar)₃ (**2**) by one-electron oxidation of [N(*n*-Bu)₄][**2**].⁶ In the synthesis of [N(*n*-Bu)₄][**2**], the tris(pentafluorophenyl)boron fragment serves the role of protecting group by preventing the formation of polymetallic μ-nitrido products, thus facilitating isolation of the first examples of molecular monometallic uranium nitrides. Herein, we report on complexes derived from the action of unprotected azide ion on **1**, a series of bimetallic uranium nitride complexes possessing a linear U=N=U core that is stable across several charge states.

Treatment of a purple-black solution of **1** in THF with NaN₃ (0.5 equiv) resulted in a gradual color change to red-brown over ~12 h. A rust-brown powder was isolated in 70% yield following workup. The paramagnetically broadened ¹H NMR spectrum, solution magnetic susceptibility ($\mu_{\text{eff}} = 3.23 \mu_{\text{B}}$, 20 °C, THF-*d*₈), and combustion analysis of the isolated material were consistent with the formation of the bimetallic diuranium(IV/IV) complex Na[(μ -N)(U(N[*t*-Bu]Ar)₃)₂] (Na[**3**]). Similarly, treatment of **1** with [N(*n*-Bu)₄][N₃] (0.5 equiv) in thawing THF provided [N(*n*-Bu)₄][**3**] ($\mu_{\text{eff}} = 3.22 \mu_{\text{B}}$, 20 °C, THF-*d*₈) in 79% yield.

With a formal 5f² electron count for each uranium(IV) center in the bimetallic [**3**][−] ion, the possibility for rich redox chemistry was realized (Scheme 1). Indeed, addition of AgOTf (1 equiv) to a solution of Na[**3**] in THF resulted in formation of the neutral diuranium(IV/V) complex (μ -N)[U(N[*t*-Bu]Ar)₃]₂ (**3**; $\mu_{\text{eff}} = 3.85 \mu_{\text{B}}$, 20 °C, chloroform-*d*), which was isolated as a brown powder in 70% yield by precipitating and filtering off the product following removal of precipitated Ag⁰ and separation from NaOTf. Furthermore, **3** reacted with [Cp₂Fe][B(Ar^F)₄] [Ar^F = 3,5-(CF₃)₂C₆H₃] in Et₂O to form the cationic diuranium(V/V) complex [(μ -N)(U(N[*t*-Bu]Ar)₃)₂][B(Ar^F)₄] ([**3**][B(Ar^F)₄]; $\mu_{\text{eff}} = 2.86 \mu_{\text{B}}$, 20 °C, chloroform-*d*), which was separated from the ferrocene coproduct by precipitation and isolated in 93% yield by filtration. The electrochemical potentials relating [**3**][−], **3**, and [**3**]⁺ were measured via cyclic voltammetry using a solution of **3** in THF. Two reversible electrochemical events at $E_{1/2} = -1.69$ and -0.64 V (vs [Cp₂Fe]^{0/+}) were observed and assigned to the [**3**][−]/**3** and **3**/[**3**]⁺ couples, respectively (Figure 1). The large separation between the [**3**][−]/**3** and **3**/[**3**]⁺ couples ($|\Delta E_{1/2}| = 1.05$ V) corresponds to an equilibrium constant $K_c \approx 5.6 \times 10^{17}$ for the comproportionation of [**3**][−] and [**3**]⁺, suggesting strong electronic communication between the uranium centers.⁷ A third electrochemical event observed at potentials above 0 V was not fully reversible at sweep rates as fast as 1000 mV s^{−1} and may be due to formation of an unstable [**3**]²⁺ species.





Figure 1. Cyclic voltammogram of **3** in THF (200 mV s^{−1} sweep rate, ~0.1 M [N(*n*-Bu)₄][B(C₆F₅)₄] supporting electrolyte) showing the two one-electron redox couples that interconvert [**3**]^{*n*} (*n* = −1, 0, +1).



Scheme 1.

The solid-state structures of $[\text{N}(n\text{-Bu})_4][\mathbf{3}]$, **3** (Figure 2), and $[\mathbf{3}][\text{B}(\text{Ar}^{\text{F}})_4]$ were determined by single-crystal X-ray diffraction, providing a rare opportunity to compare structural parameters of a U–N multiple bond over three charge states. Key metrical parameters describing the structures of $[\mathbf{3}]^n$ ($n = -1, 0, +1$) are summarized in Table 1. In all cases, each uranium center is coordinated in a trigonal pseudo- C_3 fashion by three anilide ligands and occupies a terminus of a linear $\text{U}=\text{N}=\text{U}$ bridge. Successive one-electron oxidation of the $\text{U}=\text{N}=\text{U}$ core results in a decrease of the U–N_{nitride} distance across a range of ~ 0.03 Å, while the average U–N[t-Bu]Ar distance decreases across a comparably larger range of ~ 0.12 Å. The observed contraction of the $\text{U}=\text{N}=\text{U}$ core upon oxidation is likely primarily electrostatic in origin, where the removal of metal-based nonbonding electrons results in an increasingly electron-deficient $\text{U}=\text{N}=\text{U}$ core. This structural dependence on charge is reminiscent of that displayed by penta- and hexavalent uranyl complexes, where the very covalent multiple bonds along the $\text{O}=\text{U}=\text{O}$ axis are less responsive to the charge state than are the more ionic bonds in the molecular equator.⁸ The U–N_{nitride} distances of the $[\mathbf{3}]^n$ series fall in a range that is ~ 0.1 Å longer than typical U–N distances observed in uranium imido complexes and are similar to those found in the octanuclear mixed

azide/nitride clusters $[(C_5Me_4R)_2U(\mu-N)(C_5Me_4R)U(\mu-N_3)]_4$ ($R = H, Me$) reported by Evans and co-workers^{2c} (see Table S.2 in the Supporting Information for a more detailed comparison of relevant bond metrics).





Figure 2. ORTEP rendering of **3** with ellipsoids displayed at 50% probability; hydrogen atoms and one molecule of cocrystallized Et₂O have been omitted for clarity.

Table 1. Internuclear Distances (Å) and Angles (deg) for $[(\mu-N)(U(N[t-Bu]Ar)_3)_2]^n$ ($n = -1, 0, +1$)

	[N(<i>n</i>-Bu)₄][3]	3	[3][B(Ar^F)₄]
U–N	2.080(4)	2.0625(2)	2.0470(3)
	2.077(4)		2.0511(3)
U–N[<i>t</i> -Bu]Ar (avg)	2.323	2.243	2.191
U–N–U	175.1(2)	180	180

Treatment of **[3][B(Ar^F)₄]** in THF with NaCN results in elimination of Na[B(Ar^F)₄] and formation of the known diuranium(IV/IV) cyanoimide complex $(\mu-NCN)(U(N[t-Bu]Ar)_3)_2$ (**4**), which by this new reaction was isolated in 60% yield (Scheme 2). Prior to its synthesis from NaCN and **[3][B(Ar^F)₄]**, **4** had been prepared from **1** and a cyanonitrene ($N\equiv C-N$) group transfer reagent.⁹ Here, cyanide serves as a two-electron reducing agent, converting two uranium(V) centers to uranium(IV) while inserting into the nitride bridge. This is similar to a reaction reported by Meyer and co-workers¹⁰ wherein a monometallic uranium(V) imido complex **[(L)U(NSiMe₃)]** reacts with methyl isocyanide to form Si₂Me₆ and the uranium(IV) carbodiimide derivative **[(L)U(NCNMe)]**. More generally, Scheme 2 relates to the known reactivity of certain electrophilic metal nitride complexes toward reducing Lewis bases such as phosphines or carbon monoxide, yielding

phosphiniminato or isocyanate derivatives, respectively,¹¹ and shows as well that the U=N=U core of [3]⁺ behaves as a masked metallonitrene.¹²



Scheme 2.

The complexes reported here provide insight into the structural and electrochemical characteristics of uranium nitrides. Insertion of cyanide into the U=N=U core of [3][B(Ar^F)₄] shows that the reaction chemistry of uranium nitrides extends beyond simple outer-sphere redox processes. This, along with the high polarity expected of a terminal uranium–nitride bond (U^{δ+}–N^{δ-}) and the oxidizing nature of uranium in its higher oxidation states, intimates that a high-valent *terminal* uranium nitride would have the capacity to engage in bond-forming reactions at the nitride ligand via multiple mechanistic pathways. Further developments stemming from this system of uranium nitrides may be anticipated, especially regarding reactions that result in heterolytic fragmentation of the U=N=U core.

References

- [1] (a) Fox, A. R., Bart, S. C., Meyer, K. and Cummins, C. C. *Nature* **2008**, 455, 341; (b) Silva, G. W. C., Yeaman, C. B., Sattelberger, A. P., Hartmann, T., Cerefice, G. S. and Czerwinski, K. R. *Inorg. Chem.* **2009**, 48, 10635.
- [2] (a) Fortier, S., Wu, G. and Hayton, T. W. *Dalton Trans.* **2009**, 2010; (b) Korobkov, I., Gambarotta, S. and Yap, G. P. A. *Angew. Chem., Int. Ed.* **2002**, 41, 3433; (c) Evans, W. J., Kozimor, S. A. and Ziller, J. W. *Science* **2005**, 309, 1835; (d) Evans, W. J., Miller, K. A., Ziller, J. W. and Greaves, J. *Inorg. Chem.* **2007**, 46, 8008; (e) Nocton, G., Pécaut, J. and Mazzanti, M. *Angew. Chem., Int. Ed.* **2008**, 47, 3040.
- [3] (a) Green, D. W. and Reedy, G. T. *J. Chem. Phys.* **1976**, 65, 2921; (b) Hunt, R. D., Yustein, J. T. and Andrews, L. *J. Chem. Phys.* **1993**, 98, 6070; (c) Kushto, G. P., Souter, P. F. and Andrews, L. *J. Chem. Phys.* **1998**, 108, 7121; (d) Andrews, L., Wang, X., Lindh, R., Roos, B. O. and Marsden, C. J. *Angew. Chem., Int. Ed.* **2008**, 47, 5366.
- [4] Kozimor, S. A., Yang, P., Batista, E. R., Boland, K. S., Burns, C. J., Clark, D. L., Conradson, S. D., Martin, R. L., Wilkerson, M. P. and Wolfsberg, L. E. *J. Am. Chem. Soc.* **2009**, 131, 12125.
- [5] (a) Graves, C. R., Yang, P., Kozimor, S. A., Vaughn, A. E., Clark, D. L., Conradson, S. D., Schelter, E. J., Scott, B. L., Thompson, J. D., Hay, P. J., Morris, D. E. and Kiplinger, J. L. *J. Am. Chem. Soc.* **2008**, 130, 5272; (b) Spencer, L. P., Schelter, E. J., Yang, P., Gdula, R. L., Scott, B. L., Thompson, J. D., Kiplinger, J. L., Batista, E. R. and Boncella, J. M. *Angew. Chem., Int. Ed.* **2009**, 48, 3795.
- [6] Fox, A. R. and Cummins, C. C. *J. Am. Chem. Soc.* **2009**, 131, 5716.
- [7] (a) Richardson, D. E. and Taube, H. *Inorg. Chem.* **1981**, 20, 1278; (b) Schelter, E. J., Wu, R., Scott, B. L., Thompson, J. D., Morris, D. E. and Kiplinger, J. L. *Angew. Chem., Int. Ed.* **2008**, 47, 2993.
- [8] Arnold, P. L., Love, J. B. and Patel, D. *Coord. Chem. Rev.* **2009**, 253, 1973.
- [9] Mindiola, D. J., Tsai, Y.-C., Hara, R., Chen, Q., Meyer, K. and Cummins, C. C. *Chem. Commun.* **2001**, 125.
- [10] Castro-Rodriguez, I., Nakai, H. and Meyer, K. *Angew. Chem., Int. Ed.* **2006**, 45, 2389.
- [11] Nugent, W. A. and Mayer, J. M. *Metal–Ligand Multiple Bonds*; Wiley: New York, **1988**; pp240–246.
- [12] Mindiola, D. J. *Angew. Chem., Int. Ed.* **2008**, 47, 1557.